




Adjusted puestow technique for backsliding pancreatitis-An image article

Elizabeth Swan

Open Access Publishers, 40 Bloomsbury Way, Lower Ground Floor, United Kingdom

FIGURE 1. Stamped expanded pancreatic channel and provincial liquid assortment B. Performed altered Puestow procedure. Constant backsliding (RP) pancreatitis is uncommon in youth yet can have genuine outcomes. RP drives fibrosis which makes dynamic endocrine and exocrine brokenness of pancreas. The fundamental pathologies can be inherited pancreatitis, post awful stenosis and innate formative peculiarities, for example, pancreatic divisum, pancreaticobiliary pipe oddities. A suitable ductal waste system is essential for alleviation of indications [1-4].



References

Bollen TL. Acute pancreatitis: international classification and nomenclature. *Clin Radiol.* 71, 121-133 (2016).

Lerch MM, Gorelick FS. Models of acute and chronic pancreatitis. *Gastroenterology.* 144, 1180-1193 (2013).

Thoeni RF. Imaging of Acute Pancreatitis. *Radiol Clin North Am.* 53, 1189-11208 (2015).

Tenner S, Baillie J, DeWitt J, et al. American College of Gastroenterology guideline: management of acute pancreatitis. *Am J Gastroenterol.* 108, 1400-1415 (2013).