



Comparative effectiveness of catheter ablation for atrial fibrillation: moving with the rhythm

“In order to understand the advantages and disadvantages of comparative effectiveness research, the patient, the physician, healthcare professionals, and administrators and policy makers alike must delve further into the data – the devil is in the details.”

“All scientific work is incomplete – whether it be observational or experimental. All scientific work is liable to be upset or modified by advancing knowledge. That does not confer upon us the freedom to ignore the knowledge we already have or postpone the action that it appears to demand at a given time.”

– Austin Bradford Hill (1897–1991)


On February 17, 2009, President Obama signed into law the American Recovery and Reinvestment Act. Notably, US\$1.1 billion was allotted for comparative effectiveness research (CER). On June 30, 2009, in response to the Congressional directive, the Institute of Medicine released a list of 100 initial priority topics for CER [1]. Within the first quartile, the Institute recommended an initiative “to compare the effectiveness of treatment strategies for atrial fibrillation including surgery, catheter ablation, and pharmacologic treatment”. Since atrial fibrillation (AF) is the most common arrhythmia in clinical practice, affecting 2.2 million people in the USA [2], has an estimated prevalence of up to 1% in the general population and up to 8% in those older than 80 years of age [3] and is associated with decreased survival [4], it is not a surprise that the wheels were already in motion. A systematic review of the literature, supported by the Agency for Healthcare Research and Quality (AHRQ), US Department of Health and Human Services, was well underway, and a manuscript based on the results was published on August 4, 2009 in the *Annals of Internal Medicine* [5] (the original full report is available at [101]). In summary, the systematic review found that radiofrequency ablation (RFA) of AF via a catheter-based approach after a failed drug course maintained sinus rhythm more often than continuation of drug therapy alone for up to 1 year of follow-up. In a subsequent publication of two systematic reviews (which

used different study selection criteria from the AHRQ review) of RFA and antiarrhythmic drug therapy, Calkins *et al.* also concluded that studies of RFA for AF reported higher efficacy rates than studies of antiarrhythmic drug therapy [6].

These reviews contribute to our understanding of the effectiveness of treatment modalities for AF. However, the contribution lies not only in what was found but also in what was not found about the effectiveness of treatment strategies for AF. The conclusion, if isolated, is an oversimplification of an extraordinarily complicated issue. The conclusion, if understood in its greater context, is very useful. In order to understand the advantages and disadvantages of CER, the patient, the physician, healthcare professionals, and administrators and policy makers alike must delve further into the data – the devil is in the details, as we say. Who was enrolled in the studies? As importantly, if not more so, who was excluded from the studies? What were the patient characteristics? What procedure was performed? Where and by whom was it performed? What was considered the end point, and how was it defined and ascertained? In asking and then answering these questions, it is clear that CER is important, but it is only as robust and revealing as the primary data. The conclusions apply to the patient population in the studies, which may or may not mimic ‘the real world’.

“The conclusions apply to the patient population in the studies, which may or may not mimic ‘the real world’.”

In ‘the real world’, RFA is one of the tools in the armamentarium of the cardiac electrophysiologist for the treatment of AF. At present, the indications for catheter ablation include symptomatic AF refractory or intolerant to at least one antiarrhythmic agent, first-line



Ann C Garlitski

Author for correspondence:
New England Cardiac
Arrhythmia Center,
Tufts Medical Center,
800 Washington Street,
Boston, MA 02111, USA
Tel.: +1 617 636 5902
Fax: +1 617 636 4586
agarlitski@
tuftsmedicalcenter.org

Alawi A Alsheikh-Ali

Sheikh Khalifa Medical City,
Abu Dhabi, United Arab
Emirates

Stanley Ip

*Institute for Clinical Research
and Health Policy Studies,
Tufts Medical Center, Boston,
MA, USA*

therapy in rare clinical situations and selected symptomatic AF with heart failure and/or reduced ejection fraction [7,8]. The procedure is performed in patients with paroxysmal, persistent or longstanding persistent AF. Patients may have lone AF, or they may have structural heart disease and comorbidities including ischemic heart disease, valvular heart disease, congenital heart disease, diabetes mellitus, hypertension, obesity and sleep apnea, to name a few. In other words, the patient characteristics are extremely heterogeneous. Terasawa *et al.* acknowledge that the majority of patients in the analysis had a preserved left ventricular ejection fraction, only mildly dilated atria and a mean age of 55 years, reflecting a homogeneous rather than a heterogeneous patient population [5]. Since AF is more prevalent in older patients and is more frequent in patients with mitral valve disease, heart failure, ischemia, hypertension [9] and who are likely to have reduced ejection fractions and dilated atria, there is a substantial portion of the afflicted population in whom RFA of AF has not been rigorously studied. If the primary data is not available, then CER cannot address the issue either.

.....

“The heterogeneity of the patient population is paralleled by the heterogeneity of the procedure itself. Although there is consensus that the foundation of AF ablation involves pulmonary vein isolation, there is more than one way to skin a cat.”

.....

The heterogeneity of the patient population is paralleled by the heterogeneity of the procedure itself. Although there is consensus that the foundation of AF ablation involves pulmonary vein isolation, there is more than one way to skin a cat. The procedure has evolved from ostial pulmonary vein isolation to wide area circumferential ablation to hybrid techniques, which may include additional lesion sets at the roof, the mitral isthmus, the cavotricuspid isthmus, the superior vena cava and in the coronary sinus, particularly in the case of persistent or longstanding persistent AF. Thus, it is challenging to control for all of these variations in technique, and a head-to-head comparison of antiarrhythmic drug therapy to RFA is not as simple as it may seem.

As such, while registry data may not be as methodologically rigorous as a randomized controlled trial, and despite the inherent

introduction of uncontrolled variables, this may supplement our knowledge base. In a worldwide survey of practices of RFA of AF, the rate of a major complication was 6% [10], which was relatively consistent with a major complication rate of less than 5% in most of the 84 studies in the analysis published in the *Annals of Internal Medicine* [5]. Observational data that reflects the population of patients who are actively being treated may also be hypothesis-generating and help guide the next randomized controlled trial.

.....

“Different energy sources, energy delivery systems, imaging modalities and percutaneous epicardial approaches are not on the horizon, they are here, and their effectiveness needs to be addressed.”

.....

The medical community has seen the technique of catheter ablation of AF change quickly over little more than a decade with advancements in the understanding of the pathophysiology of AF, from triggers to substrate. At even greater speed, technological innovations in the field of cardiac electrophysiology have changed the way in which the procedure is performed. Different energy sources, energy delivery systems, imaging modalities and percutaneous epicardial approaches are not on the horizon, they are here, and their effectiveness needs to be addressed. These advances are outpacing the available data at lightning speed. In order to have high-quality CER for AF, the medical community must strive to continue to gather information systematically and to perform high-quality randomized clinical trials of catheter-based ablation of AF in a wide variety of patients with long-term, well-defined end points and follow-up.

Financial & competing interests disclosure

Ann Garlitski, Alawi Alsheikh-Ali and Stanley Ip are coauthors of “Systematic review: comparative effectiveness of radiofrequency catheter ablation for atrial fibrillation. Ann. Intern. Med. 151, 191–202 (2009)”. Ann Garlitski and Alawi Alsheikh-Ali perform radiofrequency catheter ablation for atrial fibrillation. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Bibliography

- 1 Institute of Medicine: Initial national priorities for comparative effectiveness research. *Institute of Medicine*. Washington, DC, USA (2009).
- 2 Thom T, Haase N, Rosamond W *et al.*: Heart disease and stroke statistics – 2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 113(6), E85–E151 (2006).
- 3 Feinberg WM, Blackshear JL, Laupacis A, Kronmal R, Hart RG: Prevalence, age distribution, and gender of patients with atrial fibrillation. Analysis and implications. *Arch. Intern. Med.* 155(5), 469–473 (1995).
- 4 Stewart S, Hart CL, Hole DJ, McMurray JJ: A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am. J. Med.* 113(5), 359–364 (2002).
- 5 Terasawa T, Balk EM, Chung M *et al.*: Systematic review: comparative effectiveness of radiofrequency catheter ablation for atrial fibrillation. *Ann. Intern. Med.* 151(3), 191–202 (2009).
- 6 Calkins H, Reynolds MR, Spector P *et al.*: Treatment of atrial fibrillation with antiarrhythmic drugs or radiofrequency ablation: two systematic literature reviews and meta-analyses. *Circ. Arrhythmia Electrophysiol.* 2, 349–361 (2009).
- 7 Calkins H, Brugada J, Packer DL *et al.*: HRS/EHRA/ECAS expert Consensus Statement on catheter and surgical ablation of atrial fibrillation: recommendations for personnel, policy, procedures and follow-up. A report of the Heart Rhythm Society (HRS) Task Force on catheter and surgical ablation of atrial fibrillation. *Heart Rhythm* 4(6), 816–861 (2007).
- 8 Fuster V, Ryden LE, Cannom DS *et al.*: ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Revise the 2001 Guidelines for the Management of Patients With Atrial Fibrillation): developed in collaboration with the European Heart Rhythm Association and the Heart Rhythm Society. *Circulation* 114(7), E257–E354 (2006).
- 9 Levy S: Factors predisposing to the development of atrial fibrillation. *Pacing Clin. Electrophysiol.* 20(10 Pt 2), 2670–2674 (1997).
- 10 Cappato R, Calkins H, Chen SA *et al.*: Worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circulation* 111(9), 1100–1105 (2005).

Website

- 101 Agency for Healthcare Research and Quality. Comparative effectiveness review. Comparative effectiveness of radiofrequency catheter ablation for atrial fibrillation http://effectivehealthcare.ahrq.gov/repFiles/2009_0623RadiofrequencyFinal.pdf